Treatment Intervention Advisory Committee Review and Determination

Date:	April 29, 2016		
To:	DHS/DLTC ACK		
From:	Wisconsin Department of Health Services, Treatment Intervention Advisory Committee: La Collet-Klingenberg, Ph.D. (chairperson)	ına	
RE:	Determination of Art Therapy as a proven and effective treatment for individuals with autism spectrum disorder and/or other developmental disabilities	m	
☐ This is an initial review			
This is a re-review, previously reviewed in July 2013, July 2014 and April 2015.			

Section One: Overview and Determination

Please find below a statement of our determination as to whether or not the committee views Art Therapy as a proven and effective treatment for children with autism spectrum disorder and/or other developmental disabilities. In subsequent sections you will find documentation of our review process including a description of the proposed treatment, a synopsis of review findings, the treatment review evidence checklist, and a listing of the literature considered. In reviewing treatments presented to us by DHS/DLTC, we implement a review process that carefully and fully considers all available information regarding a proposed treatment. Our determination is limited to a statement regarding how established a practice is in regard to quality research. We do not make funding decisions.

Description of proposed treatment

According to a letter dated November 26, 2014 from Dr. Deaver (President) and Dr. Betts (President-elect) on behalf of the American Art Therapy Association to the TIAC, "Art therapy is a distinct mental health and behavioral science discipline that combines knowledge of human development, psychological theories and counseling techniques with training in visual arts and the creative process to provide a unique approach for helping clients improve psychological health, cognitive abilities, and sensory-motor functions. The art therapist uses art media, and often the verbal processing of produced imagery, to help people resolve conflicts and problems, develop interpersonal skills, manage behavior, reduce stress, manage pain, improve school performance, increase selfesteem, and achieve insight."

Synopsis of review

In the case of art therapy, please refer to the attached reference listing that details the reviewed research. This is the third re-review of art therapy. Below is information regarding the initial review and first re-review followed by the current re-review.

The initial review of art therapy in 2013 concluded that:

- 1. Much of the published literature involves clinical case studies or anecdotal reports that lack experimental control;
- 2. Much of the literature provides vague descriptions of relevant participant characteristics and ambiguous descriptions of treatment conditions;

- 3. Most studies fail to report concurrent treatments or combine treatments preventing examination of which treatment led to changes;
- 4. Very few studies include a control group;
- 5. The overall number of participants in empirical research involving art therapy is small;
- 6. Several studies reported art therapy is either not effective at changing the reported dependent variable or is no more effective than a control condition or other treatment; leading to a recommended Level 4-Insufficient Evidence efficacy rating.

The re-review conducted in 2014 noted one study had been published since the initial review, which was a case study. Therefore, the recommendation was that art therapy remain at a Level 4 efficacy rating.

While conducting a literature search for this review (2016), two systematic reviews were found looking at art therapy for trauma care and for care of those with cancer. Both reviews found positive outcomes and the authors encouraged the field to conduct additional research. The references for these reviews are included with the literature citations at the end of this document.

In sum, it is the decision of the committee retains an efficacy rating of Level 4- Insufficient Evidence.

Section Two: Rationale for Focus on Research Specific to Comprehensive Treatment Packages (CTP) or Models

In the professional literature, there are two classifications of interventions for individuals with Autism Spectrum Disorder (National Research Council, 2001; Odom et al., 2003; Rogers & Vismara, 2008):

- (a) **Focused intervention techniques** are individual practices or strategies (such as positive reinforcement) designed to produce a specific behavioral or developmental outcome, and
- (b) **Comprehensive treatment models** are "packages" or programs that consist of a set of practices or multiple techniques designed to achieve a broader learning or developmental impact.

To determine whether a treatment package is proven and effective, the Treatment Intervention Advisory Committee (TIAC) will adopt the following perspective as recommended by Odom et al. (2010):

The individual, focused intervention techniques that make up a comprehensive treatment model may be evidence-based. The research supporting the effectiveness of separate, individual components, however, does *not* constitute an evaluation of the comprehensive treatment model or "package." The TIAC will consider and review only research that has evaluated the efficacy of implementing the comprehensive treatment *as a package*. Such packages are most often identifiable in the literature by a consistently used name or label.

- National Research Council. (2001). *Educating children with autism*. Washington, DC: National Academy Press.
- Odom, S. L., Brown, W. H., Frey, T., Karusu, N., Smith-Carter, L., & Strain, P. (2003) Evidence-based practices for young children with autism: Evidence from single-subject research design. *Focus on Autism and Other Developmental Disabilities*, 18, 176-181.
- Odom, S. L., Boyd, B. A., Hall, L. J., & Hume, K. (2010). Evaluation of comprehensive treatment models for individuals with Autism Spectrum Disorders. *Journal of Autism and Developmental Disorders*, 40, 425-436.
- Rogers, S., & Vismara, L. (2008). Evidence-based comprehensive treatments for early autism. *Journal of Clinical Child and Adolescent Psychology*, *37*, 8-38.

Section Three: DLTC-TIAC Treatment Review Evidence Checklist

Name of Treatment: Art Therapy Level 1- Well Established or Strong Evidence (DHS 107 - Proven & Effective Treatment) Other authoritative bodies that have conducted extensive literature reviews of related treatments (e.g., National Standards Project, National Professional Development Center) have approved of or rated the treatment package as having a strong evidence base; authorities are in agreement about the level of evidence. There exist ample high quality studies that demonstrate experimental control and favorable outcomes of treatment package. Minimum of two group studies or five single subject studies or a combination of the two. Studies were conducted across at least two independent research groups. Studies were published in peer reviewed journals. There is a published procedures manual for the treatment, or treatment implementation is clearly defined (i.e., replicable) within the studies. Participants (i.e., N) are clearly identified as individuals with autism spectrum disorders or developmental disabilities. Notes: At this level, include ages of participants and disabilities identified in body of research Level 2 – Established or Moderate Evidence (DHS 107 - Proven & Effective Treatment) Other authoritative bodies that have conducted extensive literature reviews of related treatments (e.g., National Standards Project, NPDC) have approved of or rated the treatment package as having at least a minimal evidence base; authorities may not be in agreement about the level of evidence. There exist at least two high quality studies that demonstrate experimental control and favorable outcomes of treatment package. ☐ Minimum of one group study or two single subject studies or a combination of the two. ☐ Studies were conducted by someone other than the creator/provider of the treatment. ☐ Studies were published in peer reviewed journals. Participants (i.e., N) are clearly identified as individuals with autism spectrum disorders or developmental disabilities.

Notes: At this level, include ages of participants and disabilities identified in body of research

<u>Level 3 – Emerging Evidence (DHS 107 – Promising as a Proven & Effective Treatment)</u>		
 Other authoritative bodies that have conducted extensive literature reviews of related treatments (e.g., National Standards Project, NPDC) have recognized the treatment package as having an emerging evidence base; authorities may not be in agreement about the level of evidence. There exists at least one high quality study that demonstrates experimental control and favorable outcomes of treatment package. May be one group study or single subject study. Study was conducted by someone other than the creator/provider of the treatment. Study was published in peer reviewed journal. Participants (i.e., N) are clearly identified as individuals with autism spectrum disorders or developmental disabilities. 		
de veropmentar disubilities.		
Notes: At this level, include ages of participants and disabilities identified in body of research		
<u>Level 4 – Insufficient Evidence (Experimental Treatment)</u>		
Other authoritative bodies that have conducted extensive literature reviews of related treatments (e.g., National Standards Project, NPDC) have not recognized the treatment package as having an emerging evidence base; authorities are in agreement about the level of evidence.		
There is not at least one high quality study that demonstrates experimental control and favorable		
outcomes of treatment package.		
 Study was conducted by the creator/provider of the treatment. Study was not published in a peer reviewed journal. 		
Participants (i.e., N) are not clearly identified as individuals with autism spectrum disorders or developmental disabilities.		
Notes:		

p. 5

<u>Level 5 – Untested (Experimental Treatment) &/or Potentially Harmful</u>	
 Other authoritative bodies that have conducted extensive literature reviews of related treatments (e.g., National Standards Project, NPDC) have not recognized the treatment package as having an emerging evidence base; authorities are in agreement about the level of evidence. There are no published studies supporting the proposed treatment package. 	
 There exists evidence that the treatment package is potentially harmful. Authoritative bodies have expressed concern regarding safety/outcomes. Professional bodies (i.e., organizations or certifying bodies) have created statements regarding safety/outcomes. 	
<i>Notes</i> : At this level, please specify if the treatment is reported to be potentially harmful, providing documentation	

Committee Members Completing Initial Review of Research Base: Lana Collet-Klingenberg, Shannon Stuart

Committee Decision on Level of Evidence to Suggest the Proposed Treatment is Proven and Effective: Level 4- Insufficient Evidence

References Supporting Identification of Evidence Levels:

Date: April 29, 2016

- Chambless, D.L., Hollon, S.D. (1998). Defining empirically supported therapies. *Journal of Consulting and Clinical Psychology*, 66(1) 7-18.
- Chorpita, B.F. (2003). The frontier of evidence---based practice. In A.E. Kazdin & J.R. Weisz (Eds.). *Evidence-based psychotherapies for children and adolescents* (pp. 42---59). New York: The Guilford Press.
- Odom, S. L., Collet-Klingenberg, L., Rogers, S. J., & Hatton, D. (2010). Evidence-based practices in interventions for children and youth with autism spectrum disorders. *Preventing School Failure*, 54(4), 275-282.

Section Four: Literature Review

Cite all literature reviewed here and note month of most recent article reviewed for future reviewers:

No new articles found since April 2015.

Review articles found referencing art therapy for trauma care and for those with cancer diagnoses:

- Archer, S., Buxton, S., & Sheffield, D. (2015). The effect of creative psychological interventions on psychological outcomes for adult cancer patients: A systematic review of randomised controlled trials. *Psycho-Oncology*, 24, 1-10. ***systematic review
- Schouten, K.A., DeNiet, G.J., Knipscheer, J.W., Kleber, R.J., & Hutschemaekers, G.J.M. (2015) The effectiveness of art therapy in the treatment of traumatized adults: A systematic review on art therapy and trauma. *Trauma, Violence, & Abuse, 16(2), 220-228.*

Journal articles provided by the Wisconsin Art Therapy Association:

- Emery, M. J. (2004). Art therapy as an intervention for autism. Art Therapy: *Journal of the American Art Therapy Association*, 21(3), 143-147. ***case study
- Epp, K. M. (2008). Outcome-based evaluation of a social skills program using art therapy and group therapy for children on the autism spectrum. *Children and Schools*, 30(1), 27-36. ***pre/post design
- Hess, K. L., Morrier, M. J., Heflin, L. J., & Ivey, M. L. (2008). Autism treatment survey: Services received by children with autism spectrum disorders in public school classrooms. *Journal of Autism and Developmental Disorders*, 38, 961-971. ***survey study
- Martin, N. (2009). Art therapy and autism: Overview and recommendations. Art Therapy: *Journal of the American Art Therapy Association*, 26(4), 187-190. ***recommendations paper
- Martin, N. (2008). Assessing portrait drawings created by children and adolescents with an autism spectrum disorder. Art Therapy: *Journal of the American Art therapy Association*, 25(1), 15-23. ***pilot study
- Simpson, R. L. (2005). Evidence-based practices and students with autism spectrum disorders. *Focus on Autism and Other Developmental Disabilities*, 20(3), 140-149. ***literature review

Journal articles reviewed in the 2014 review:

Durani, H. (2014). Facilitating attachment in children with autism through art therapy: A case study. *Journal of Psychotherapy Integration*, 24(2), 99-108. ***case study

Literature reviewed in the initial review (2013):

- Banks, S., David, P., Howard, V.F., & McLaughlin, T.F. (1993). The effects of directed art activities on the behavior of young children with disabilities: A multielement baseline analysis. Art Therapy: *Journal of the American Art Therapy Association*, 10, 235-240.
- Chapman, L., Morabita, D., Ladakakos, C., Schreier, H., & Knudson, M.M. (2001). The effectiveness of art therapy interventions in reducing post traumatic stress disorder (PTSD) symptoms in pediatric trauma patients. Art Therapy: *Journal of the American Art Therapy Association*, 18, 100-104.

- Coholic, D., Eys, M., & Lougheed, S. (2012). Investigating the effectiveness of an artsbased and mindfulness-based group program for the improvement of resilience in children in need. *Journal of Child Family Studies*, 21, 833-844.
- Favara-Scacco, C., Smirne, G., Schiliro, G., & Di Cataldo, A. (2001). Art therapy as support for children with leukemia during painful procedures. Medical and Pediatric Oncology, 36, 474-480.
- Kearns, D. (2004). Art therapy with a child experiencing sensory integration difficulty. Art Therapy: *Journal of the American Art Therapy Association*, 21, 95-101.
- Lyshak-Stelzer, F., Singer, P., St. John, P., & Chemtob, C.M. (2007). Art therapy for adolescents with posttraumatic stress disorder symptoms: A pilot study. Art Therapy: *Journal of the American Art Therapy Association*, 24, 163-169.
- Regev, D., & Guttmann, J. (2005). The psychological benefits of artwork: The case of children with learning disorders. *The Arts in Psychotherapy*, *32*, 302-312.
- Reynolds, M.W., Nabors, L., & Quinlan, A. (2000). The effectiveness of art therapy: Does it work? Art Therapy: *Journal of the American Art Therapy Association*, 17, 207-213.
- Sanders, E. (2003). "Looks aren't everything": Pediatric spinal cord injuries and art therapy. Trauma and Loss: *Research and Interventions*, *3*, 31-36.
- Slayton, S.C., D'Archer, J., & Kaplan, F. (2010). Outcome studies on the efficacy of art therapy: A review of the findings. Art Therapy: *Journal of the American Art Therapy Association*, 27, 108-118.
- Stafstrom, C.E., Havlena, J., & Krezinski, A.J. (2012). Art therapy focus groups for children and adolescents with epilepsy. *Epilepsy & Behavior*, 24, 227-233.
- Viscardi, N. (1994). Art therapy as a support group for adolescents with muscular dystrophy. American *Journal of Art Therapy*, *32*, 66.